

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) (PMEL/1825R), 1mg/mL, Clone: [PMEL/1825R], Rabbit, Monoclonal BOT-BNUM1825-50

Artikelname	gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) (PMEL/1825R), 1mg/mL, Clone: [PMEL/1825R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1825-50
Hersteller Artikelnummer	BNUM1825-50
Alternativnummer	BOT-BNUM1825-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human SILV protein
Produktbeschreibung	Cytotoxic T lymphocytes (CTLs) recognize melanoma-associated antigens, which belong to three main groups. These groups include tumor-associated testis-specific antigens, melanocyte differentiation antigens and mutated or aberrantly expressed antigens...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[PMEL/1825R]
Molekulargewicht	90-100 kDa

UniProt	P40967
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user