

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, pan (Epithelial Marker), 1mg/mL, Rabbit BOT-BNUM1835-50

Artikelname	Cytokeratin, pan (Epithelial Marker), 1mg/mL, Rabbit
Artikelnummer	BOT-BNUM1835-50
Hersteller Artikelnummer	BNUM1835-50
Alternativnummer	BOT-BNUM1835-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	KRT77 proteins Recombinant full-length human KRT76
Produktbeschreibung	Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, which 67 kDa (CK1), 64 kDa (CK3), 59 kDa (...)
Konzentration	1 mg/mL
Molekulargewicht	40-67 kDa
UniProt	Q01546 , Q7Z794
Puffer	PBS, no BSA, no azide
Quelle	Animal

Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.25-0.5 ug/mL Immunofluorescence 1-2 ug/mL Western blotting 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user
------------------------	--