

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) (PCRP-CDX2-1A3), 1 mg/mL, IgG1, Clone: [PCRP-CDX2-1A3], Mouse, Monoclonal BOT-BNUM1920-50**

Artikelname	CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) (PCRP-CDX2-1A3), 1 mg/mL, IgG1, Clone: [PCRP-CDX2-1A3], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1920-50
Hersteller Artikelnummer	BNUM1920-50
Alternativnummer	BOT-BNUM1920-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human CDX2protein
Produktbeschreibung	The specificity of this monoclonal antibody to its intended target was validated by HuProt(TM)Array, containing more than 19,000, full-length human proteins.The intestine-specific transcription factors CDX1 and CDX2 are important for directing intest...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[PCRP-CDX2-1A3]
Molekulargewicht	40 kDa

Isotyp	IgG1
UniProt	<a href="#">Q99626</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Immunoprecipitation 2-4 ug/mg protein Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user