

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

GP2 (Glycoprotein 2) / ZAP75 (GP2/2569R), 1mg/mL, Clone: [GP2/2569R], Rabbit, Monoclonal BOT-BNUM2569-50

| | |
|--------------------------|---|
| Artikelname | GP2 (Glycoprotein 2) / ZAP75 (GP2/2569R), 1mg/mL, Clone: [GP2/2569R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUM2569-50 |
| Hersteller Artikelnummer | BNUM2569-50 |
| Alternativnummer | BOT-BNUM2569-50-50UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human GP2 protein fragment (around aa35-179) (exact sequence is proprietary) |
| Produktbeschreibung | GP2 (glycoprotein 2), also known as ZAP75, is a 537 amino acid secreted protein. It is an integral membrane protein that is secreted from intracellular zymogen granules and associates with the plasma membrane via glycosylphosphatidylinositol (GPI) li... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [GP2/2569R] |

| | |
|------------------------|--|
| Molekulargewicht | 59 kDa |
| UniProt | P55259 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |