

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), 1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal BOT-BNUM2651-50

Artikelname	Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), 1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2651-50
Hersteller Artikelnummer	BNUM2651-50
Alternativnummer	BOT-BNUM2651-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human Interleukin 10 (IL10) protein
Produktbeschreibung	Interleukin 10 (IL10) is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immuno-regulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II an...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[IL10/2651R]
Molekulargewicht	20 kDa (Monomer), 37 kDa (Dimer)

UniProt	P22301
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user