

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TCL1 (T-Cell Marker) (TCL1/2747R), 1mg/mL, Clone: [TCL1/2747R], Rabbit, Monoclonal BOT-BNUM2747-50

Artikelname	TCL1 (T-Cell Marker) (TCL1/2747R), 1mg/mL, Clone: [TCL1/2747R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2747-50
Hersteller Artikelnummer	BNUM2747-50
Alternativnummer	BOT-BNUM2747-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human TCL1 protein fragment (around aa 2-109) (exact sequence is proprietary)
Produktbeschreibung	T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromos...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[TCL1/2747R]

Molekulargewicht	14 kDa
UniProt	P56279
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user