

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AMACR / p504S (Prostate Cancer Marker) (AMACR/2748R), 1mg/mL, Clone: [AMACR/2748R], Rabbit, Monoclonal BOT-BNUM2748-50

Artikelname	AMACR / p504S (Prostate Cancer Marker) (AMACR/2748R), 1mg/mL, Clone: [AMACR/2748R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2748-50
Hersteller Artikelnummer	BNUM2748-50
Alternativnummer	BOT-BNUM2748-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human AMACR protein fragment (around aa 297-394) (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of 42 kDa, which is identified as Alpha-methylacyl-CoA Racemase (AMACR), also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and betbeta-oxidation of branched-chain fatty acids. AMACR is ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[AMACR/2748R]

Molekulargewicht	42 kDa
UniProt	Q9UHK6
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user