

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 18 (KRT18) (KRT18/2808R), 1mg/mL, Clone: [KRT18/2808R], Rabbit, Monoclonal BOT-BNUM2808-50**

Artikelname	Cytokeratin 18 (KRT18) (KRT18/2808R), 1mg/mL, Clone: [KRT18/2808R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2808-50
Hersteller Artikelnummer	BNUM2808-50
Alternativnummer	BOT-BNUM2808-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human KRT18 protein
Produktbeschreibung	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[KRT18/2808R]
Molekulargewicht	45 kDa

UniProt	<a href="#">P05783</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western Blot 0.5-2 ug/mL Optimal dilution for a specific application should be determined by user