

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Golgi Complex (Marker for Human Cells) (GLG1/2829R), 1mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal BOT-BNUM2829-50

Artikelname	Golgi Complex (Marker for Human Cells) (GLG1/2829R), 1mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2829-50
Hersteller Artikelnummer	BNUM2829-50
Alternativnummer	BOT-BNUM2829-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	SU-DHL-1 large cell lymphoma cells.
Produktbeschreibung	This MAb recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[GLG1/2829R]
Molekulargewicht	Not Known

UniProt	Not Known
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user