

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Tenascin C (Stromal Marker For Epithelial Malignancy) (TNC/2981R), 1mg/mL, Clone: [TNC/2981R], Rabbit, Monoclonal BOT-BNUM2981-50

Artikelname	Tenascin C (Stromal Marker For Epithelial Malignancy) (TNC/2981R), 1mg/mL, Clone: [TNC/2981R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2981-50
Hersteller Artikelnummer	BNUM2981-50
Alternativnummer	BOT-BNUM2981-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human TNC protein
Produktbeschreibung	In Western blotting, this antibody reacts with two bands of ~MW of 210 kDa and 300 kDa, identified as two isoforms of Tenascin C. Specificity of this MAb is validated by sequential immunoprecipitation with a PAb against Tenascin C. Tenascin C is a mu...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[TNC/2981R]

Molekulargewicht	210 kDa and 300 kDa
UniProt	P24821
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with rat Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user