

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

LMO2 (B-Cell Marker)(LMO2/3147R), CF740 conjugate, 0.1mg/mL, Clone: [LMO2/3147R], CF 740, Rabbit, Monoclonal BOT-BNC743147-100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | LMO2 (B-Cell Marker)(LMO2/3147R), CF740 conjugate, 0.1mg/mL, Clone: [LMO2/3147R], CF 740, Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC743147-100 |
| Hersteller Artikelnummer | BNC743147-100 |
| Alternativnummer | BOT-BNC743147-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 23-140) human LMO2 protein (exact sequence is proprietary) |
| Konjugation | CF 740 |
| Produktbeschreibung | The LMO2 protein has a central and crucial role in hematopoietic development and is highly conserved. It has a particular function in normal and lymphatic endothelial cells involving the regulation of angiogenesis and lymph-angiogenesis. Immunohistoc... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Klon-Bezeichnung | [LMO2/3147R] |
| Molekulargewicht | 24 kDa |
| UniProt | P25791 |
| Puffer | PBS, 0.1% rBSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Flow cytometry: 1-2 ug/million cells, Immunofluorescence: 1-2 ug/mL, Optimal dilution for a specific application should be determined. Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody |