

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, pan (Epithelial Marker)(PCK/3150), CF740 conjugate, 0.1mg/mL, Clone: [PCK/3150], CF 740, Mouse, Monoclonal BOT-BNC743150-100

| | |
|--------------------------|---|
| Artikelname | Cytokeratin, pan (Epithelial Marker)(PCK/3150), CF740 conjugate, 0.1mg/mL, Clone: [PCK/3150], CF 740, Mouse, Monoclonal |
| Artikelnummer | BOT-BNC743150-100 |
| Hersteller Artikelnummer | BNC743150-100 |
| Alternativnummer | BOT-BNC743150-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Canine, Feline, Human |
| Immunogen | Crude cytokeratin extract prepared from RT-4 and MCF-7 cells. |
| Konjugation | CF 740 |
| Produktbeschreibung | Anti-cytokeratin clone PCK/3150 demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, PCK/3150 is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast ducts, bronchial ep... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [PCK/3150] |

| | |
|------------------------|--|
| Molekulargewicht | 40-67 kDa (Multiple) |
| UniProt | Multiple |
| Puffer | PBS, 0.1% rBSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |