

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human IL2 protein, His tag (active), Unconjugated GTX00265-PRO

Artikelname	Human IL2 protein, His tag (active), Unconjugated
Artikelnummer	GTX00265-PRO
Hersteller Artikelnummer	GTX00265-pro
Alternativnummer	GTX00265-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	3558
UniProt	P60568
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ala21~Thr153 (NP_000577.2)

Anwendungsbeschreibung	Interleukin-2 (IL-2) is an interleukin, a type of cytokine signaling molecule in the immune system. IL-2 has essential roles in key functions of the immune system, tolerance and immunity, primarily via its direct effects on T cells. In the thymus, where T cells mature, it prevents autoimmune diseases by promoting the differentiation of certain immature T cells into regulatory T cells, which suppress other T cells that are otherwise primed to attack normal healthy cells in the body. IL-2 also promotes the differentiation of T cells into effector T cells and into memory T cells when the initial T cell is also stimulated by an antigen, thus helping the body fight off infections. To measure the effect of IL-2 on cell proliferation, CTLL-2 cells were seeded into triplicate wells of 96-well plates at a density of 2000 cells/well with 2% serum standard RPMI-1640 contain various concentrations of recombinant human IL-2. After incubated for 96h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37C. Proliferation of CTLL-2 cells after incubation with IL-2 for 96h observed by inverted microscope.
------------------------	---