

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant Human TNF RII (C-mFc) EBT-EPT211

Artikelname	Recombinant Human TNF RII (C-mFc)
Artikelnummer	EBT-EPT211
Hersteller Artikelnummer	EPT211
Alternativnummer	EBT-EPT211-50
Hersteller	ELK Biotechnology
Kategorie	Proteine/Peptide
Produktbeschreibung	Recombinant Human Tumor Necrosis Factor Receptor Superfamily Member 1B is produced by our Mammalian expression system and the target gene encoding Pro24-Thr206 is expressed with a mFc tag at the C-terminus....
Molekulargewicht	Molecular weight: 46.44 KDa. Apparent molecular weight: 60 KDa, reducing conditions
UniProt	P20333
Reinheit	Greater than 95% as determined by reducing SDS-PAGE.

Anwendungsbeschreibung	Redissolve: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.. Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test. Background: Tumor necrosis factor receptor superfamily member 1B (TNFRSF1B) is a member of the tumor necrosis factor receptor superfamily. Human TNF RII contains four cysteine-rich repeats in its ECD, which shares 58% and 56% amino acid sequence identity with the mouse and rat orthologs, respectively. TNF RII is expressed predominantly on cells of the hematopoietic lineage, such as T and natural killer cells, as well as on endothelial cells, microglia, astrocytes, neurons, oligodendrocytes, cardiac myocytes, thymocytes, and mesenchymal stem cells. TNF RII binds to the membrane-bound forms of TNFα and Lymphotoxinα/TNFβ; soluble TNF is thought to signal predominantly through TNF RI. Soluble TNF RII is believed to inhibit TNF biological activity by binding TNF thereby preventing it from activating membrane TNF receptors
------------------------	---