

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-U2AF65/U2AF2 Antibody Picoband, Rabbit, Polyclonal BOB-A03639-2-CARRIER-FREE

|                          |  |
|--------------------------|--|
| Artikelname              | Anti-U2AF65/U2AF2 Antibody Picoband, Rabbit, Polyclonal  |
| Artikelnummer            | BOB-A03639-2-CARRIER-FREE  |
| Hersteller Artikelnummer | A03639-2-carrier-free  |
| Alternativnummer         | BOB-A03639-2-CARRIER-FREE-100UG  |
| Hersteller               | Boster Bio   |
| Wirt                     | Rabbit   |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, FC, ICC, IF, IHC, IP, WB  |
| Spezies Reaktivität      | Human, Mouse, Rat  |
| Immunogen                | E.coli-derived human U2AF65/U2AF2 recombinant protein (Position: M238-H470).   |
| Produktbeschreibung      | Boster Bio Anti-U2AF65/U2AF2 Antibody Picoband catalog A03639-2. Tested in ELISA, Flow Cytometry, IP, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees... |
| Klonalität               | Polyclonal   |
| Konzentration            | Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.  |
| Molekulargewicht         | Observed Molecular Weight: 65 kDa. Calculated Molecular Weight: 28461 MW   |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">11338</a>  |
| UniProt                | <a href="#">P26368</a>   |
| Puffer                 | Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .   |
| Reinheit               | Immunogen affinity purified.   |
| Formulierung           | Lyophilized  |
| Target-Kategorie       | Splicing factor U2AF 65 kDa subunit  |
| Application Verdünnung | Western blot, 0.25-0.5µg/ml, Human, Mouse, Rat<br>Immunohistochemistry (Paraffin-embedded Section), 0.5-1µg/ml, Human, Mouse, Rat<br>Immunocytochemistry/Immunofluorescence, 2µg/ml, Human<br>Immunoprecipitation, 0.5-2 µg/ml, Human<br>Flow Cytometry (Fixed), 1-3µg/1x10 <sup>6</sup> |