

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Monoclonal Antibody to CD15 / FUT4 (Reed-Sternberg Cell Marker)(Clone : BRA-4F1), Clone: [BRA-4F1], Mouse ABI-36-1162

| | |
|--------------------------|---|
| Artikelname | Monoclonal Antibody to CD15 / FUT4 (Reed-Sternberg Cell Marker)(Clone : BRA-4F1), Clone: [BRA-4F1], Mouse |
| Artikelnummer | ABI-36-1162 |
| Hersteller Artikelnummer | 36-1162 |
| Alternativnummer | ABI-36-1162-100UG |
| Hersteller | Abeomics |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Myelomonocytic leukemia cells |
| Produktbeschreibung | CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on >95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, CD15 is expressed in Reed-Sternberg cell... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [BRA-4F1] |
| NCBI | 2526 |
| UniProt | P22083 |

| | |
|------------------------|---|
| Reinheit | Affinity Chromatography |
| Formulierung | Purified |
| Target-Kategorie | CD15 / FUT4 (Reed-Sternberg Cell Marker) |
| Application Verdünnung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA Buffer, pH 7.5-8.5, for 45 min at 95 |