

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Monoclonal Antibody to Androgen Receptor (Marker of Androgen Dependence)(Clone : DHTR/882), Clone: [DHTR/882], Mouse ABI-36-1318

|                          |   |
|--------------------------|---|
| Artikelname              | Monoclonal Antibody to Androgen Receptor (Marker of Androgen Dependence)(Clone : DHTR/882), Clone: [DHTR/882], Mouse  |
| Artikelnummer            | ABI-36-1318   |
| Hersteller Artikelnummer | 36-1318   |
| Alternativnummer         | ABI-36-1318-100UG   |
| Hersteller               | Abeomics  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FACS, IF, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human DHTR protein  |
| Produktbeschreibung      | Recognizes a protein of 110kDa, which is identified as androgen receptor (AR). It reacts with full length, and the newly described A form of the receptor. It does not cross react with estrogen, progesterone, or glucocorticoid receptors. The expressio... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [DHTR/882]  |
| NCBI                     | <a href="#">367</a>   |
| UniProt                  | <a href="#">P10275</a>  |

|                        |   |
|------------------------|---|
| Reinheit               | Affinity Chromatography   |
| Formulierung           | Purified  |
| Target-Kategorie       | Androgen Receptor (Marker of Androgen Dependence)   |
| Application Verdünnung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min |