

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Monoclonal Antibody to NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker)(Clone : SPM299), Clone: [SPM299], Mouse ABI-36-1516

| | |
|--------------------------|---|
| Artikelname | Monoclonal Antibody to NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker)(Clone : SPM299), Clone: [SPM299], Mouse |
| Artikelnummer | ABI-36-1516 |
| Hersteller Artikelnummer | 36-1516 |
| Alternativnummer | ABI-36-1516-100UG |
| Hersteller | Abeomics |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | NGFR from A875 melanoma cells |
| Produktbeschreibung | It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). Its epitope spans in aa 1-160 of extracellular domain of NGFR/NTR. NGF-receptor contains an extracellula... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM299] |
| NCBI | 4804 |
| UniProt | P08138 |

| | |
|------------------------|---|
| Reinheit | Affinity Chromatography |
| Formulierung | Purified |
| Target-Kategorie | NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) |
| Application Verdünnung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min a |