

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Monoclonal Antibody to Bcl-2 (Apoptosis & Follicular Lymphoma Marker)(BCL2/782 + BCL2/796), Clone: [BCL2/782 + BCL2/796], Mouse ABI-36-1647

Artikelname	Monoclonal Antibody to Bcl-2 (Apoptosis & Follicular Lymphoma Marker)(BCL2/782 + BCL2/796), Clone: [BCL2/782 + BCL2/796], Mouse
Artikelnummer	ABI-36-1647
Hersteller Artikelnummer	36-1647
Alternativnummer	ABI-36-1647-100UG
Hersteller	Abeomics
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human bcl-2 protein
Produktbeschreibung	This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 i oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 i oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphoma...
Klonalität	Monoclonal
Klon-Bezeichnung	[BCL2/782 + BCL2/796]
NCBI	596

UniProt	P10415
Reinheit	Affinity Chromatography
Formulierung	Purified
Target-Kategorie	Bcl-2 (Apoptosis & Follicular Lymphoma Marker)
Application Verdünnung	Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95°C followed by cooling at RT for 20 minutes)