

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Monoclonal Antibody to Complement 4d (C4d) (Acute Humoral Rejection Marker)(Clone : C4D204), Clone: [C4D204], Mouse ABI-36-1767

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Monoclonal Antibody to Complement 4d (C4d) (Acute Humoral Rejection Marker)(Clone : C4D204), Clone: [C4D204], Mouse |
| Artikelnummer | ABI-36-1767 |
| Hersteller Artikelnummer | 36-1767 |
| Alternativnummer | ABI-36-1767-100UG |
| Hersteller | Abeomics |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Complement 4d protein |
| Produktbeschreibung | This MAb is specific to Complement 4d (C4d) and it reacts with the secreted as well as cell-bound C4d. C4d is a degradation product of the activated complement factor C4b. Complement 4b is typically activated by binding of Abs to specific target mole... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [C4D204] |
| NCBI | 720 |
| UniProt | P0C0L4 |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reinheit | Affinity Chromatography |
| Formulierung | Purified |
| Target-Kategorie | Complement 4d (C4d) (Acute Humoral Rejection Marker) |
| Application Verdünnung | ELISA (For coating use Ab at 1-2ug/ml order Ab without BSA), Immunohistochemistry (Formalin-fixed) (1:200-1:400 for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95°C fo |