

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Monoclonal Antibody to von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(Clone : IIIE2.34), Clone: [IIIE2.34], Mouse ABI-36-1807

| | |
|--------------------------|---|
| Artikelname | Monoclonal Antibody to von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(Clone : IIIE2.34), Clone: [IIIE2.34], Mouse |
| Artikelnummer | ABI-36-1807 |
| Hersteller Artikelnummer | 36-1807 |
| Alternativnummer | ABI-36-1807-100UG |
| Hersteller | Abeomics |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC, IP, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human vWF fragment spanning aa 845-949 |
| Produktbeschreibung | von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [IIIE2.34] |
| NCBI | 7450 |
| UniProt | P04275 |

| | |
|------------------------|---|
| Reinheit | Affinity Chromatography |
| Formulierung | Purified |
| Target-Kategorie | von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) |
| Application Verdünnung | Immunoprecipitation (0.5-1 g/500ug protein lysate), Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues re |