

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Donkey IgG anti-Chicken IgG (H+L)-FITC, MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh DNA-SEC-182806

Artikelname	Donkey IgG anti-Chicken IgG (H+L)-FITC, MinX Bo,Go,Gp,Hm,Ho,Hu,Ms,Rb,Rt,Sh
Artikelnummer	DNA-SEC-182806
Hersteller Artikelnummer	SEC-182806
Alternativnummer	DNA-SEC-182806
Hersteller	dianova
Wirt	Donkey
Kategorie	Antikörper
Applikation	FLISA,FACS,IF
Spezies Reaktivität	Gallus
Immunogen	Chicken IgG whole molecule
Konjugation	FITC
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Goat,Guinea pig,Hamster (all),Equine,Human,Mouse,Rabbit,Rat,Sheep

Produktbeschreibung	Anti-Chicken IgG Fluorescein Antibody generated in donkey detects chicken IgY. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria,...
Klonalität	Polyclonal
Konzentration	1 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by conjugation to fluorochrome and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Donkey Serum, Chicken IgG and Chicken Serum. No reaction was observed against Bovine, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit, Rat and Sheep Serum Proteins. This antibody will react with heavy chains of Chicken IgG and with light chains of most Chicken immunoglobulins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Chicken
Antibody Type	Secondary Antibody
Application Verdünnung	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.0, IF Microscopy Dilution: 1:1,000 - 1:5,000
Anwendungsbeschreibung	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.