

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Mouse IgG (H+L)-FITC, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Rb,Rt,Sh DNA-SEC-183128

Artikelname	Goat IgG anti-Mouse IgG (H+L)-FITC, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Rb,Rt,Sh
Artikelnummer	DNA-SEC-183128
Hersteller Artikelnummer	SEC-183128
Alternativnummer	DNA-SEC-183128
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	FLISA,FACS,IHC,IF,WB
Spezies Reaktivität	Mouse
Immunogen	Mouse IgG whole molecule
Konjugation	FITC
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Bovine,Gallus,Goat,Guinea pig,Hamster (all),Equine,Human,Rabbit,Rat,Sheep

Produktbeschreibung	Anti-Mouse IgG Fluorescein Antibody generated in goat detects reactivity to Mouse IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, b...
Klonalität	Polyclonal
Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Goat Anti-Mouse IgG was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Mouse
Antibody Type	Secondary Antibody
Application Verdünnung	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.08, Immunohistochemistry Dilution: User Optimized, IF Microscopy Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: User Optimized
Anwendungsbeschreibung	Anti-Mouse IgG Fluorescein Antibody has been tested by dot blot and western blot and is ideal for Fluorescent Western Blot, FLISA, Flow Cytometry, Immunohistochemistry and Immunofluorescence Microscopy as well as other antibody detection methods.