

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Rabbit IgG (Fc)-HRPO, MinX none DNA-SEC-183365

Artikelname	Goat IgG anti-Rabbit IgG (Fc)-HRPO, MinX none
Artikelnummer	DNA-SEC-183365
Hersteller Artikelnummer	SEC-183365
Alternativnummer	DNA-SEC-183365
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Rabbit
Immunogen	Anti-Rabbit IgG F(c) fragment was produced by repeated immunization with Rabbit IgG F(c) fragment in goat.
Konjugation	HRPO
Format	IgG
Spezifität	IgG (Fc)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Rabbit IgG F(c) peroxidase antibody generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion...

Klonalität	Polyclonal
Konzentration	2.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Anti-Rabbit IgG F(c) antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab).
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target-Kategorie	Rabbit
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:110,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:20,000
Anwendungsbeschreibung	Anti-Rabbit IgG Fc antibody is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.