

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Hamster armenian IgG (H+L)-unconj., MinX none DNA-SEC-183643

Artikelname	Goat IgG anti-Hamster armenian IgG (H+L)-unconj., MinX none
Artikelnummer	DNA-SEC-183643
Hersteller Artikelnummer	SEC-183643
Alternativnummer	DNA-SEC-183643
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,ChIP,IHC,IP,WB
Spezies Reaktivität	Hamster (armenian)
Immunogen	Armenian Hamster IgG whole molecule
Konjugation	Unconjugated
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Armenian Hamster IgG Antibody generated in goat detects Armenian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacte...
Klonalität	Polyclonal

Konzentration	5.03 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Armenian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Armenian Hamster IgG and Armenian Hamster Serum. Reactivity with Golden Syrian Hamster IgG is greatly reduced.
Formulierung	Liquid (sterile filtered)
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target-Kategorie	Armenian Hamster
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:40,000, ChIP Dilution: User Optimized, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Immunoprecipitation Dilution: User Optimized, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-Armenian Hamster IgG Antibody has been tested by ELISA and is suitable for immunoprecipitation, immunodiffusion, conjugation and most immunological methods requiring lot-to-lot consistency, high titer and specificity.