

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Hamster (all) IgG (H+L)-unconj., MinX Ms,Rt DNA-SEC-183648

Artikelname	Goat IgG anti-Hamster (all) IgG (H+L)-unconj., MinX Ms,Rt
Artikelnummer	DNA-SEC-183648
Hersteller Artikelnummer	SEC-183648
Alternativnummer	DNA-SEC-183648
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Hamster (all)
Immunogen	Armenian and Golden Syrian Hamster IgG, whole molecule
Konjugation	Unconjugated
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	Mouse,Rat
Produktbeschreibung	Anti-Golden Syrian & Armenian Hamster IgG Antibody generated in goat detects Golden Syrian & Armenian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunogl...

Klonalität	Polyclonal
Konzentration	1.158
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Secondary Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian and Armenian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Golden Syrian Hamster IgG and Armenian Hamster IgG. No reaction was observed against Mouse or Rat serum proteins. Specificity was confirmed using ELISA at less than 1% cross reactivity.
Formulierung	Liquid (sterile filtered)
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target-Kategorie	Armenian and Golden Syrian Hamster
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Anwendungsbeschreibung	Anti-Golden Syrian & Armenian Hamster IgG antibody has been tested by ELISA and dot blot and is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.