

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat IgG anti-Hamster (all) IgG (H+L)-Biotin, MinX none DNA-SEC-183652

Artikelname	Goat IgG anti-Hamster (all) IgG (H+L)-Biotin, MinX none
Artikelnummer	DNA-SEC-183652
Hersteller Artikelnummer	SEC-183652
Alternativnummer	DNA-SEC-183652
Hersteller	dianova
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA,IHC,WB
Spezies Reaktivität	Hamster (all)
Immunogen	Armenian and Golden Syrian Hamster IgG, whole molecule
Konjugation	Biotin
Format	IgG
Spezifität	IgG (H+L)
Minimale Kreuzreaktivität (MinX)	no cross-adsorbtion
Produktbeschreibung	Anti-Golden Syrian & Armenian Hamster IgG Biotin Antibody generated in goat detects Golden Syrian & Armenian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. I...
Klonalität	Polyclonal

Konzentration	1.0 mg/mL
Isotyp	Ig
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reinheit	Anti-Golden Syrian and Aramenian hamster IgG was prepared from monospecific antiserum by immunoaffinity chromatography using Armenian/Golden Syrian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Armenian/Golden Syrian Hamster IgG and Armenian/Golden Syrian Hamster Serum. No reaction was observed against Mouse or Rat Serum Proteins.
Formulierung	Lyophilized
Formel	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target-Kategorie	Armenian and Golden Syrian Hamster
Antibody Type	Secondary Antibody
Application Verdünnung	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,000, Western Blot Dilution: 1:1,000 - 1:5,000
Anwendungsbeschreibung	Anti Golden Syrian & Armenian Hamster IgG Biotin conjugated antibody has been tested by ELISA and western blot and is suitable for Immunohistochemistry. Specific conditions for reactivity should be optimized by the end user.