

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Murine CXCL16 Antibody, Goat, Polyclonal ABT-ABG10058-U100

| | |
|----------------------------|---|
| Article Name | Anti-Murine CXCL16 Antibody, Goat, Polyclonal |
| Biozol Catalog Number | ABT-ABG10058-U100 |
| Supplier Catalog Number | ABG10058-U100 |
| Alternative Catalog Number | ABT-ABG10058-U100-100UG |
| Manufacturer | Abcepta |
| Host | Goat |
| Category | Antikörper |
| Application | ELISA, WB |
| Species Reactivity | Mouse |
| Clonality | Polyclonal |
| Purity | Produced from sera of goats pre-immunized with highly pure (>98%) recombinant mCXCL16. Anti-Murine CXCL16 specific antibody was purified by affinity chromatography employing immobilized mCXCL16 matrix. |
| Form | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2. |
| Antibody Type | Polyclonal Antibody |

Application Notes

WesternBlot: To detect mCXCL16 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mCXCL16 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect mCXCL16 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Murine CXCL16 (61-062BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mCXCL16.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.