

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Rat IFN-gamma Antibody, Goat, Polyclonal ABT-ABG10165-U100

Article Name	Anti-Rat IFN-gamma Antibody, Goat, Polyclonal
Biozol Catalog Number	ABT-ABG10165-U100
Supplier Catalog Number	ABG10165-U100
Alternative Catalog Number	ABT-ABG10165-U100-100UG
Manufacturer	Abcepta
Host	Goat
Category	Antikörper
Application	ELISA, WB
Species Reactivity	Rat
Clonality	Polyclonal
Purity	Produced from sera of goats pre-immunized with highly pure recombinant Rat IFN-gamma. Anti-Rat IFN-gamma specific antibody was purified by affinity chromatography employing immobilized Rat IFN-gamma matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect Rat IFN-gamma by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Rat IFN-gamma is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect Rat IFN-gamma by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Rat IFN-gamma (62-061GBT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IFN-gamma.. Neutralization: To yield one-half maximal inhibition [ND<span style=font-size: 16px,>50] of the biological activity of Rat IFN-gamma (0.3 ng/ml), a concentration of 60.0-65.0 ng/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.