

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Human IGF-BP7 Antibody, Rabbit, Polyclonal ABT-ABG10180-U100

Article Name	Anti-Human IGF-BP7 Antibody, Rabbit, Polyclonal
Biozol Catalog Number	ABT-ABG10180-U100
Supplier Catalog Number	ABG10180-U100
Alternative Catalog Number	ABT-ABG10180-U100-100UG
Manufacturer	Abcepta
Host	Rabbit
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Human
Clonality	Polyclonal
Purity	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIGF-BP7. Anti-Human IGF-BP7 specific antibody was purified by affinity chromatography employing immobilized hIGF-BP7 matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect hIGF-BP7 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIGF-BP7 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hIGF-BP7 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IGF-BP7 (60-194BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIGF-BP7.. Immunohistochemistry: This antibody stained HeLa and U-251 MG cells. The primary antibody was incubated at 2.0 mg/ml overnight at 4C followed by a fluorescent labeled secondary antibody. Optimal concentrations and conditions may vary. Information and photo are courtesy of the Cell Profiling group, SciLifeLab Stockholm.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.