

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human IL-11 Antibody, Rabbit, Polyclonal ABT-ABG10197-U050

Article Name	Anti-Human IL-11 Antibody, Rabbit, Polyclonal
Biozol Catalog Number	ABT-ABG10197-U050
Supplier Catalog Number	ABG10197-U050
Alternative Catalog Number	ABT-ABG10197-U050-50UG
Manufacturer	Abcepta
Host	Rabbit
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Human
Clonality	Polyclonal
Purity	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-11. Anti-Human IL-11 specific antibody was purified by affinity chromatography employing immobilized hIL-11 matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect hIL-11 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-11 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hIL-11 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IL-11 (60-011BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-11.. Immunohistochemistry: This antibody stained human brain stroke (including control cortex and stroke core areas) tissue. The primary antibody was incubated at 2.5 mg/ml overnight at 4°C. This was followed by a fluorophore conjugated secondary antibody. Optimal concentrations and conditions may vary.

 Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm.. Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hIL-11 (1.50 ng/ml), a concentration of 0.02-0.05 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.