

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human IL-6 Antibody, Mouse, Monoclonal ABT-ABG10300-U500

Article Name	Anti-Human IL-6 Antibody, Mouse, Monoclonal
Biozol Catalog Number	ABT-ABG10300-U500
Supplier Catalog Number	ABG10300-U500
Alternative Catalog Number	ABT-ABG10300-U500-500UG
Manufacturer	Abcepta
Host	Mouse
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Human
Clonality	Monoclonal
Purity	Produced in BALB/c mice using highly pure (>98%) recombinant human IL-6 as the immunizing antigen. This IgG1K antibody was purified from cell culture by Protein A affinity chromatography.
Form	A sterile filtered antibody solution was lyophilized from PBS.
Antibody Type	Monoclonal Antibody

Application Notes

WesternBlot: To detect hIL-6 by Western Blot analysis this antibody can be used at a concentration of 0.20-0.40 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-6 is 0.5-1.0 ng/lane, under reducing or non-reducing conditions.. Sandwich: In a sandwich ELISA (assuming 100µl/well), a concentration of 4.0-8.0 µg/ml of this antibody will detect at least 500 pg/ml of recombinant human IL-6 when used with BioGems biotinylated antigen affinity purified anti-human IL-6 (60-006BT) as the detection antibody at a concentration of approximately 0.5-1.0 µg/ml.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human cervical squamous cell carcinoma. The recommended concentration is 0.125 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary. Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute.

 *Additional Immunostaining data available. Please contact Tech Support for information.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.