

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human IL-8 Antibody, Rabbit, Polyclonal ABT-ABG10308-U100

Article Name	Anti-Human IL-8 Antibody, Rabbit, Polyclonal
Biozol Catalog Number	ABT-ABG10308-U100
Supplier Catalog Number	ABG10308-U100
Alternative Catalog Number	ABT-ABG10308-U100-100UG
Manufacturer	Abcepta
Host	Rabbit
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Human
Clonality	Polyclonal
Purity	Produced from sera of rabbits pre-immunized with highly pure recombinant Human IL-8. Anti-Human IL-8 specific antibody was purified by affinity chromatography employing immobilized Human IL-8 matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect Human IL-8 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human IL-8 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect Human IL-8 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IL-8 (60-008BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human IL-8.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of normal human skin. The recommended concentrations are 1.0 µg/ml for two hours at room temperature or 0.25 µg/ml overnight at 4C. An HRP-labeled polymer detection system was used with a non-alcohol soluble AEC chromogen and a proteinase K antigen retrieval. Optimal concentrations and conditions may vary.
 *Additional Immunostaining data available. Please contact Tech Support for information.. Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Human IL-8 (100ng/ml), a concentration of 2.0 - 3.0 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.