

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Murine IP-10 Antibody, Rabbit, Polyclonal ABT-ABG10316-U100

Article Name	Anti-Murine IP-10 Antibody, Rabbit, Polyclonal
Biozol Catalog Number	ABT-ABG10316-U100
Supplier Catalog Number	ABG10316-U100
Alternative Catalog Number	ABT-ABG10316-U100-100UG
Manufacturer	Abcepta
Host	Rabbit
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Mouse
Clonality	Polyclonal
Purity	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mIP-10. Anti-Murine IP-10 specific antibody was purified by affinity chromatography employing immobilized mIP-10 matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect mIP-10 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIP-10 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect mIP-10 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Murine IP-10 (61-081BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mIP-10.. Immunohistochemistry: This antibody stained colchicine injected mouse brain (including the hippocampus region) tissue. The primary antibody was incubated at 1.0 mg/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA(TM)) reagent. Optimal concentrations and conditions may vary. . Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of mIP-10 (100.0 ng/ml), a concentration of 10.0 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.