

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human MCP-2 Antibody, Rabbit, Polyclonal ABT-ABG10357-U100

Article Name	Anti-Human MCP-2 Antibody, Rabbit, Polyclonal
Biozol Catalog Number	ABT-ABG10357-U100
Supplier Catalog Number	ABG10357-U100
Alternative Catalog Number	ABT-ABG10357-U100-100UG
Manufacturer	Abcepta
Host	Rabbit
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Human
Clonality	Polyclonal
Purity	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMCP-2. Anti-Human MCP-2 specific antibody was purified by affinity chromatography employing immobilized hMCP-2 matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect hMCP-2 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMCP-2 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hMCP-2 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human MCP-2 (60-213BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMCP-2.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human normal gastric mucosa. The recommended concentration is 0.75mg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

 Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.