

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Biotinylated Anti-Human MEC Antibody, Rabbit, Polyclonal ABT-ABG10386-U025**

|                            |   |
|----------------------------|---|
| Article Name               | Biotinylated Anti-Human MEC Antibody, Rabbit, Polyclonal  |
| Biozol Catalog Number      | ABT-ABG10386-U025   |
| Supplier Catalog Number    | ABG10386-U025   |
| Alternative Catalog Number | ABT-ABG10386-U025-25UG  |
| Manufacturer               | Abcepta   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | ELISA, WB   |
| Species Reactivity         | Human   |
| Clonality                  | Polyclonal  |
| Purity                     | Produced from sera of rabbits immunized with highly pure recombinant Human MEC. Anti-Human MEC specific antibody was purified by affinity chromatography and then biotinylated. |
| Form                       | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.  |
| Antibody Type              | Polyclonal Antibody   |

Application Notes

WesternBlot: To detect hMEC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMEC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hMEC by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems Polyclonal Anti-Human MEC (60-218P) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMEC.. Direct: To detect hMEC by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMEC.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.