

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human MIG Antibody, Rabbit, Polyclonal ABT-ABG10393-U050

| | |
|----------------------------|---|
| Article Name | Anti-Human MIG Antibody, Rabbit, Polyclonal |
| Biozol Catalog Number | ABT-ABG10393-U050 |
| Supplier Catalog Number | ABG10393-U050 |
| Alternative Catalog Number | ABT-ABG10393-U050-50UG |
| Manufacturer | Abcepta |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA, IHC, WB |
| Species Reactivity | Human |
| Clonality | Polyclonal |
| Purity | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMIG. Anti-Human MIG specific antibody was purified by affinity chromatography employing immobilized hMIG matrix. |
| Form | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2. |
| Antibody Type | Polyclonal Antibody |

Application Notes

WesternBlot: To detect hMIG by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIG is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hMIG by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human MIG (60-223BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIG.. Immunohistochemistry: This antibody stained formalin-fixed paraffin-embedded sections of human renal tumor with parenchyma tissue. The recommended concentration is 0.25µg/mL- 0.50µg/mL with an overnight incubation at 4C. An HRP-labeled polymer detection system was used with a DAB chromogen. Optimal results for these conditions were achieved with heat induced antigen retrieval with a pH 6.0 sodium citrate buffer. Optimal concentrations and conditions may vary.. Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hMIG (100.0 ng/ml), a concentration of 5.0-7.0 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.