

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Steroidogenic Factor 1 (SF-1) Antibody, IgG1, Clone: [rNR5A1/4369], Mouse, Monoclonal NBT-2516-MSM9-P0

| | |
|----------------------------|---|
| Article Name | Recombinant Steroidogenic Factor 1 (SF-1) Antibody, IgG1, Clone: [rNR5A1/4369], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-2516-MSM9-P0 |
| Supplier Catalog Number | 2516-MSM9-P0 |
| Alternative Catalog Number | NBT-2516-MSM9-P0-20,NBT-2516-MSM9-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant fragment (around aa220-461) of human SF-1 protein |
| Product Description | Steroidogenic Factor 1 (SF-1) is considered an orphan nuclear receptor that belongs to subfamily 5. It was found to be a regulator of steroidogenic enzyme gene expression. Oxysterols are suggested as its ligands. It is expressed in all steroidogenic ... |
| Clonality | Monoclonal |
| Clone Designation | [rNR5A1/4369] |
| Molecular Weight | 53kDa. |
| Isotype | IgG1 |

| | |
|-------------------|--|
| NCBI | 2516 |
| UniProt | Q13285 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |