

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD15 / FUT4 (Reed-Sternberg Cell Marker) Antibody, IgM, Clone: [MY-1], Mouse, Monoclonal NBT-2526-MSM10-P1ABX

| | |
|----------------------------|---|
| Article Name | CD15 / FUT4 (Reed-Sternberg Cell Marker) Antibody, IgM, Clone: [MY-1], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-2526-MSM10-P1ABX |
| Supplier Catalog Number | 2526-MSM10-P1ABX |
| Alternative Catalog Number | NBT-2526-MSM10-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IF, IHC |
| Species Reactivity | Human |
| Immunogen | Purified neutrophils from normal human peripheral blood |
| Product Description | CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on 95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, CD15 is expressed in Reed-Sternberg cells... |
| Clonality | Monoclonal |
| Clone Designation | [MY-1] |
| Molecular Weight | 220kDa |
| Isotype | IgM |

| | |
|-------------------|--|
| NCBI | 2526 |
| UniProt | P22083 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95degC followed by cooling at RT |