

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Blood Group Antigen H Type 2 (CD173) Antibody, IgM, Clone: [A46-B/B10], Mouse, Monoclonal NBT-28-MSM2-P0

| | |
|----------------------------|---|
| Article Name | Blood Group Antigen H Type 2 (CD173) Antibody, IgM, Clone: [A46-B/B10], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-28-MSM2-P0 |
| Supplier Catalog Number | 28-MSM2-P0 |
| Alternative Catalog Number | NBT-28-MSM2-P0-20,NBT-28-MSM2-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IF |
| Species Reactivity | Human |
| Immunogen | Human breast cancer MCF-7 cells |
| Product Description | Recognizes the blood group H type 2 antigens, trisaccharide Fuc(TM)1-2GalA°a1-4GlcNAcA°a1 of human origin. The specificity of this MAb has been identified by haemagglutination tests, immunohistochemistry, binding inhibition studies, and absorption ex... |
| Clonality | Monoclonal |
| Clone Designation | [A46-B/B10] |
| Molecular Weight | Multiple |
| Isotype | IgM |

| | |
|-------------------|--|
| NCBI | 28 |
| UniProt | P16442 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Haemagglutination, Binding Inhibition Assays, Immunofluorescence (2-4ug/ml), Immunohistology (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 mi |