

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Tenascin C (Stromal Marker for Epithelial Malignancy) Antibody, IgG1, Clone: [SPM319], Mouse, Monoclonal NBT-3371-MSM1X-P0

| | |
|----------------------------|---|
| Article Name | Tenascin C (Stromal Marker for Epithelial Malignancy) Antibody, IgG1, Clone: [SPM319], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-3371-MSM1X-P0 |
| Supplier Catalog Number | 3371-MSM1X-P0 |
| Alternative Catalog Number | NBT-3371-MSM1X-P0-20,NBT-3371-MSM1X-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | Human |
| Immunogen | Human breast carcinoma |
| Product Description | In Western blotting, it reacts with two bands of ~MW of 210kDa and 300kDa, identified as two isoforms of Tenascin C. Specificity of this MAb is validated by sequential immunoprecipitation with a PAb against Tenascin C. Tenascin C is a multifunctional... |
| Clonality | Monoclonal |
| Clone Designation | [SPM319] |
| Molecular Weight | 210kDa and 300kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 3371 |
| UniProt | P24821 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 mi |