

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, IgG1, Clone: [F8/86], Mouse, Monoclonal NBT-7450-MSM6-P0

Article Name	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) Antibody, IgG1, Clone: [F8/86], Mouse, Monoclonal
Biozol Catalog Number	NBT-7450-MSM6-P0
Supplier Catalog Number	7450-MSM6-P0
Alternative Catalog Number	NBT-7450-MSM6-P0-20,NBT-7450-MSM6-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	von Willebrand Factor isolated from human plasma
Product Description	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Clonality	Monoclonal
Clone Designation	[F8/86]
Molecular Weight	250kDa
Isotype	IgG1

NCBI	7450
UniProt	P04275
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f