

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) Antibody, Clone: [KRTH/1576R], Rabbit, Monoclonal NBT-RBM22-1576-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) Antibody, Clone: [KRTH/1576R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-RBM22-1576-P1ABX |
| Supplier Catalog Number | RBM22-1576-P1ABX |
| Alternative Catalog Number | NBT-RBM22-1576-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Canine, Human |
| Immunogen | Recombinant full-length human KRT76 protein |
| Product Description | This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67kDa (CK1), 64kDa (CK3), 59kDa (CK4), 58kDa (CK5), 56kDa (CK6), 52kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subf... |
| Clonality | Monoclonal |
| Clone Designation | [KRTH/1576R] |
| Molecular Weight | 52-67kDa |
| NCBI | 51350 |

| | |
|-------------------|--|
| UniProt | Q01546 |
| Form | 200ug/ml of recombinant MAb Purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution |