

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

ODC-1 Antibody [ODC1/485], Unconjugated, Mouse, Monoclonal PRS-33-182

| | |
|----------------------------|------------------------------------------------------------------------------|
| Article Name | ODC-1 Antibody [ODC1/485], Unconjugated, Mouse, Monoclonal |
| Biozol Catalog Number | PRS-33-182 |
| Supplier Catalog Number | 33-182 |
| Alternative Catalog Number | PRS-33-182-100 |
| Manufacturer | ProSci |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant human ODC-1 protein was used as the immunogen for this antibody. |
| Conjugation | Unconjugated |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [ODC1/485] |
| NCBI | 4953 |
| UniProt | P11926 |
| Buffer | PBS with 0.1 mg/ml BSA and 0.05% sodium azide |
| Form | Liquid |

| | |
|--------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Dilute | Optimal dilutions for each application to be determined by the researcher |
| Application Notes | <p>Flow Cytometry: 0.5-1 ug/million cells IF: 0.5-1 ug/ml WB: 0.5-1 ug/ml IHC (FFPE): 0.5-1 ug/ml for 30 minutes at RT (1) Prediluted format : incubate for 30 min at RT (2)</p> <p>The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titered up or down for optimal performance.</p> <p>1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |