

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CTLA4 Antibody [2G10], IgG1, Unconjugated, Mouse, Monoclonal PRS-RF16012

| | |
|----------------------------|--|
| Article Name | CTLA4 Antibody [2G10], IgG1, Unconjugated, Mouse, Monoclonal |
| Biozol Catalog Number | PRS-RF16012 |
| Supplier Catalog Number | RF16012 |
| Alternative Catalog Number | PRS-RF16012-0.02, PRS-RF16012-0.1 |
| Manufacturer | ProSci |
| Host | Mouse |
| Category | Antikörper |
| Application | ELISA, ICC, IF, IHC-P, WB |
| Species Reactivity | Human |
| Immunogen | CTLA-4 antibody was raised against the extracellular domain of human CTLA-4. |
| Conjugation | Unconjugated |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [2G10] |
| Molecular Weight | Predicted: 25 kDa Observed: 32 kDa |
| Isotype | IgG1 |
| NCBI | 1493 |
| UniProt | P16410 |

| | |
|--------------------|--|
| Buffer | CTLA-4 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol. |
| Form | Liquid |
| Application Dilute | Optimal dilutions for each application to be determined by the researcher. |
| Application Notes | CTLA-4 antibody can be used for detection of CTLA-4 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples, Immunocytochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested. |