

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

PDL2 Antibody [4E10], IgG1, Unconjugated, Mouse, Monoclonal PRS-RF16021

| | |
|----------------------------|--|
| Article Name | PDL2 Antibody [4E10], IgG1, Unconjugated, Mouse, Monoclonal |
| Biozol Catalog Number | PRS-RF16021 |
| Supplier Catalog Number | RF16021 |
| Alternative Catalog Number | PRS-RF16021-0.02, PRS-RF16021-0.1 |
| Manufacturer | ProSci |
| Host | Mouse |
| Category | Antikörper |
| Application | ELISA, FC, ICC, IF, IHC-P, WB |
| Species Reactivity | Human |
| Immunogen | PD-L2 antibody was raised against the extracellular domain of human PD-L2. |
| Conjugation | Unconjugated |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [4E10] |
| Molecular Weight | Predicted: 30 kDa Observed: 38 kDa |
| Isotype | IgG1 |
| NCBI | 80380 |
| UniProt | P9BQ51 |

| | |
|--------------------|---|
| Buffer | PD-L2 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol. |
| Form | Liquid |
| Application Dilute | Optimal dilutions for each application to be determined by the researcher. |
| Application Notes | PD-L2 antibody can be used for detection of PD-L2 by Western blot at 0.5 - 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples, Immunocytochemistry in human samples, Immunofluorescence in human samples and Flow Cytometry in mouse samples. All other applications and species not yet tested. |