

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker)(VU-4H5), CF405S conjugate, 0.1mg/mL, Clone: [VU-4H5], Mouse, Monoclonal BOT-BNC040431-100

Article Name	MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker)(VU-4H5), CF405S conjugate, 0.1mg/mL, Clone: [VU-4H5], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC040431-100
Supplier Catalog Number	BNC040431-100
Alternative Catalog Number	BOT-BNC040431-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Synthetic glycosylated MUC1 60mer tandem repeat NH2-(HGVTSAPDT(GalNAc)RPAPGSTAPPAHG)3- COOH, conjugated to bovine serum albumin
Conjugation	CF405S
Product Description	MAB VU-4H5 reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. The dominant epitope of MAB VU4H5 is APDTR as established with epitope fingerprinting. VU-4H5 preferentially binds to under...
Clonality	Monoclonal
Concentration	0.1 mg/mL

Clone Designation	[VU-4H5]
Molecular Weight	265-400 kDa
UniProt	P15941
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user