

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Chromogranin A(LK2H10 + PHE5), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [LK2H10 PHE5], Mouse, Monoclonal BOT-BNC040698-100**

Article Name	Chromogranin A(LK2H10 + PHE5), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [LK2H10 PHE5], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC040698-100
Supplier Catalog Number	BNC040698-100
Alternative Catalog Number	BOT-BNC040698-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Monkey, Mouse, Porcine, Rat
Immunogen	Human pheochromocytoma (LK2H10 & PHE5)
Conjugation	CF405S
Product Description	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[LK2H10 PHE5]

Molecular Weight	68-75 kDa
Isotype	IgG1
UniProt	<a href="#">P10645</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user