

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker), CF405S conjugate, 0.1mg/mL, Rabbit BOT-BNC040728-500**

Article Name	IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker), CF405S conjugate, 0.1mg/mL, Rabbit
Biozol Catalog Number	BOT-BNC040728-500
Supplier Catalog Number	BNC040728-500
Alternative Catalog Number	BOT-BNC040728-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Purified human Ig Gamma Chain
Conjugation	CF405S
Product Description	This antibody recognizes a protein of 75 kDa, identified as gamma heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), mu (IgM), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes...
Concentration	0.1 mg/mL
Molecular Weight	75 kDa
UniProt	<a href="#">P01861</a> , <a href="#">P01860</a> , <a href="#">P01859</a> , <a href="#">P01857</a>

Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user