**BIOZOL**
Diagnostica Vertrieb GmbH, Oehleckerring 11-13
22419 Hamburg, Germany
**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99
**E-Mail:** info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

# Product Datasheet

## Cdc20(CDC20/1102), CF405S conjugate, 0.1mg/mL, Clone: [CDC20/1102], Mouse, Monoclonal
## BOT-BNC041102-500

| | |
|---|---|
| Article Name | Cdc20(CDC20/1102), CF405S conjugate, 0.1mg/mL, Clone: [CDC20/1102], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC041102-500 |
| Supplier Catalog Number | BNC041102-500 |
| Alternative Catalog Number | BOT-BNC041102-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human Cdc20 protein |
| Conjugation | CF405S |
| Product Description | Cyclins, regulatory subunits that associate with kinases, control many of the important steps in cell cycle progression. The Cdc2 protein kinase (p34Cdc2) exhibits protein kinase activity in vitro and exists in a complex with both cyclin B and a prot... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [CDC20/1102] |

| | |
|---|---|
| Molecular Weight | 55 kDa |
| UniProt | Q12834 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-fixed 2-4 ug/mL\|Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes\|Immunofluorescence 0.5-1 ug/mL\|Flow Cytometry 0.5-1 ug/million cells/0.1 mL\|Optimal dilution for a specific application should be determined by user |